

## **Barium Enema Preparation Instructions**

 Date:
 Time:
 Location: St. Francis Hospital 2<sup>nd</sup> Floor Suite 202

### Items to Purchase:

- □ 4 Dulcolax Laxative tablets
- □ 8.3 oz bottle of <u>Miralax</u>
- □ 10 oz bottle of Magnesium Citrate
- G4 oz of Gatorade (<u>NO</u> red or purple)

### Three (3) days prior to your procedure:

- 1. Restrict/limit your diet- <u>Do not</u> eat nuts, seeds, popcorn, or corn.
- 2. <u>Stop</u> taking any fiber supplements, such as Metamucil, Citrucel, Fiberall, etc.

# **One day prior to the procedure:** \**If you are diabetic, you must check your blood sugar before taking any medications*\*

- Clear liquids only (see below). <u>Do not</u> drink anything that is **RED** or **PURPLE**. <u>Foods allowed under Clear Liquid Diet:</u> <u>Soups:</u> Clear bouillon, broth, or consommé <u>Beverages:</u> Tea, coffee, decaffeinated tea/coffee, Kool-Aid (<u>no RED or PURPLE</u>), carbonated beverages. **DO NOT** put any milk or cream in your tea/coffee. <u>Juices:</u> White cranberry, apple, grape, strained lemonade/limeade, orange drink [any juice you can see through and without pulp is acceptable] <u>Dessert:</u> water ices, Italian ice(s), popsicles, and jello (<u>no RED or PURPLE</u>)
   At 5:00pm- Take 4 Dulcolax tablets/pills
- 3. Apply Vaseline (petroleum jelly) or Desitin cream to rectum to improve comfort before loose stools/diarrhea starts.
- 4. At 6:00pm- mix the 8.3 oz of Miralax with 64 oz of Gatorade. Shake the solution until the Miralax is dissolved. Drink a total of five (5) 8oz glasses, one glass every 10-15 minute until all solution is ingested.
- 5. Continue drinking clear liquids until bedtime.
- 6. If stools are not clear or the consistency of tea by 9:00pm, take the full bottle of magnesium and then drink 1 glass of water every 30 minutes, up to 3 glasses.
- 7. <u>\*\*NOTHING TO EAT OR DRINK AFTER MIDNIGHT (12AM)\*\*</u>

### On the day of your procedure:

- 1. Take your blood pressure and heart medications with a sip of water. No other medication until after the procedure.
- 2. Arrive at the location at least 30 minutes prior to your scheduled test time.